

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Unintentional Injuries/Violence**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN9: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle, one or more times during the 30 days before the survey)	21.0 (19.7-22.4)	3,938	28.2 (25.6-30.9)	1,189		●	
QN10: Percentage of students who drove a car or other vehicle when they had been drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	4.9 (3.9-6.1)	1,557	14.2 (11.2-17.9)	434		●	
QN12: Percentage of students who carried a weapon (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)	20.2 (18.8-21.5)	3,848	22.8 (20.3-25.4)	1,112			●
QN13: Percentage of students who carried a weapon on school property (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)	5.6 (4.9-6.4)	3,881	8.5 (6.9-10.4)	1,138		●	
QN15: Percentage of students who did not go to school because they felt unsafe at school or on their way to or from school (on at least 1 day during the 30 days before the survey)	8.1 (7.3-9.0)	3,907	15.4 (13.4-17.7)	1,164		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Unintentional Injuries/Violence**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN16: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club, one or more times during the 12 months before the survey)	9.1 (8.2-10.1)	3,939	18.1 (15.9-20.5)	1,194		●	
QN17: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)	33.6 (32.1-35.2)	3,795	40.1 (37.2-43.2)	1,118		●	
QN18: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	16.1 (14.9-17.4)	3,841	20.6 (18.2-23.2)	1,139		●	
QN19: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	10.3 (9.4-11.3)	3,886	20.1 (17.8-22.7)	1,116		●	
QN20: Percentage of students who experienced sexual violence (being forced by anyone to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey)	13.2 (12.2-14.4)	3,785	27.7 (25.0-30.5)	1,096		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

### District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN21: Percentage of students who experienced sexual dating violence (being forced by someone they were dating or going out with to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)	4.5 (3.8-5.3)	2,944	13.3 (11.1-15.9)	808		●	
QN22: Percentage of students who experienced physical dating violence (being physically hurt on purpose by someone they were dating or going out with [counting such things as being hit, slammed into something, or injured with an object or weapon] one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)	8.2 (7.2-9.2)	3,041	22.2 (19.5-25.3)	866		●	
QN23: Percentage of students who were bullied on school property (ever during the 12 months before the survey)	9.4 (8.5-10.4)	3,881	19.5 (17.2-22.0)	1,118		●	
QN24: Percentage of students who were electronically bullied (counting being bullied through texting, Instagram, Facebook, or other social media, ever during the 12 months before the survey)	8.8 (7.9-9.8)	3,870	18.3 (16.1-20.8)	1,112		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Unintentional Injuries/Violence**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN25: Percentage of students who felt sad or hopeless (almost every day for >=2 weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)	33.2 (31.7-34.7)	3,900	50.5 (47.4-53.5)	1,110		●	
QN26: Percentage of students who seriously considered attempting suicide (ever during the 12 months before the survey)	17.3 (16.1-18.5)	3,868	40.0 (37.0-43.0)	1,122		●	
QN27: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)	16.0 (14.8-17.2)	3,888	33.5 (30.7-36.5)	1,120		●	
QN28: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	13.7 (12.5-15.1)	3,203	31.4 (28.5-34.5)	985		●	
QN29: Percentage of students who had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	5.0 (4.3-5.9)	3,172	12.9 (10.8-15.4)	939		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Tobacco Use**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN31: Percentage of students who first tried cigarette smoking before age 13 years (even one or two puffs)	9.3 (8.4-10.3)	3,885	21.3 (18.9-23.9)	1,138		●	
QN32: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	4.3 (3.7-4.9)	3,885	13.3 (11.4-15.5)	1,141		●	
QNFRCIG: Percentage of students who currently smoked cigarettes frequently (on 20 or more days during the 30 days before the survey)	0.4 (0.3-0.7)	3,885	2.0 (1.3-3.1)	1,141		●	
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	0.3 (0.2-0.6)	3,885	1.5 (0.9-2.4)	1,141		●	
QN35: Percentage of students who currently used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo], on at least 1 day during the 30 days before the survey)	14.7 (13.7-15.9)	3,891	24.8 (22.4-27.4)	1,176		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Tobacco Use**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QNFREVP: Percentage of students who currently used electronic vapor products frequently (on 20 or more days during the 30 days before the survey)	1.9 (1.5-2.3)	3,891	3.3 (2.5-4.4)	1,176		●	
QNDAYEVP: Percentage of students who currently used electronic vapor products daily (on all 30 days during the 30 days before the survey)	1.1 (0.8-1.4)	3,891	2.3 (1.6-3.3)	1,176		●	
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Copenhagen, Grizzly, Skoal, or Camel Snus], not counting any electronic vapor products, on at least 1 day during the 30 days before the survey)	4.1 (3.4-4.8)	3,942	15.7 (13.7-18.1)	1,194		●	
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars, on at least 1 day during the 30 days before the survey)	5.8 (5.1-6.6)	3,932	17.1 (15.0-19.5)	1,198		●	
QNTB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)	7.8 (7.0-8.7)	3,907	21.1 (18.8-23.6)	1,185		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Tobacco Use**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QNTB3: Percentage of students who currently smoked cigarettes or cigars or used smokeless tobacco (on at least 1 day during the 30 days before the survey)	9.0 (8.2-10.0)	3,913	24.5 (22.0-27.1)	1,192		●	
QNTB4: Percentage of students who currently smoked cigarettes or cigars or used smokeless tobacco or electronic vapor products (on at least 1 day during the 30 days before the survey)	18.8 (17.6-20.0)	3,879	32.8 (30.1-35.7)	1,187		●	
QNTB5: Percentage of students who currently smoked cigarettes or used electronic vapor products (on at least 1 day during the 30 days before the survey)	16.4 (15.2-17.5)	3,864	28.2 (25.7-31.0)	1,162		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Alcohol/Other Drug Use**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN40: Percentage of students who had their first drink of alcohol before age 13 years (other than a few sips)	18.7 (17.5-20.1)	3,764	33.0 (30.1-35.9)	1,116		●	
QN41: Percentage of students who currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)	24.1 (22.8-25.5)	3,726	40.7 (37.7-43.8)	1,071		●	
QN42: Percentage of students who currently were binge drinking (had four or more drinks of alcohol in a row for female students or five or more drinks of alcohol in a row for male students, within a couple of hours, on at least 1 day during the 30 days before the survey)	9.4 (8.5-10.3)	3,830	20.5 (18.2-23.1)	1,110		●	
QN46: Percentage of students who tried marijuana for the first time before age 13 years	14.1 (12.9-15.3)	3,847	22.8 (20.4-25.5)	1,141		●	
QN47: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	39.0 (37.4-40.6)	3,882	51.3 (48.3-54.3)	1,154		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.



## 2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

### District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Alcohol/Other Drug Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN49: Percentage of students who ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)	20.9 (19.6-22.3)	3,939	37.9 (35.0-40.8)	1,192		●	
QN50: Percentage of students who ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)	4.5 (3.9-5.2)	3,904	16.1 (14.0-18.4)	1,183		●	
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	9.1 (8.2-10.1)	3,942	24.5 (22.1-27.2)	1,192		●	
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)	3.4 (2.9-4.1)	3,936	14.5 (12.5-16.7)	1,190		●	
QN53: Percentage of students who ever used methamphetamines (also called "speed," "crystal meth," "crank," "ice," or "meth," one or more times during their life)	3.3 (2.8-4.0)	3,905	13.9 (11.9-16.1)	1,182		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Alcohol/Other Drug Use**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)	4.4 (3.7-5.1)	3,940	14.5 (12.5-16.7)	1,188		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Sexual Behaviors**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN58: Percentage of students who ever had sexual intercourse	83.0 (81.8-84.2)	3,397	72.1 (69.0-75.0)	901	●		
QN59: Percentage of students who had sexual intercourse for the first time before age 13 years	14.0 (12.8-15.4)	3,404	13.1 (11.0-15.6)	937			●
QN60: Percentage of students who had sexual intercourse with four or more persons during their life	23.2 (21.7-24.7)	3,382	18.3 (15.9-21.0)	919	●		
QN61: Percentage of students who were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)	57.6 (55.8-59.3)	3,379	52.2 (48.8-55.5)	920	●		
QN62: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	15.3 (13.7-17.1)	1,867	20.6 (17.2-24.4)	462		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Sexual Behaviors**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN63: Percentage of students who did not use a condom during last sexual intercourse (among students who were currently sexually active)	39.5 (37.2-41.9)	1,813	53.0 (47.5-58.4)	347		●	
QN64: Percentage of students who did not use birth control pills before last sexual intercourse (to prevent pregnancy among students who were currently sexually active)	89.9 (88.4-91.2)	1,809	90.2 (86.3-93.1)	299			●
QNIUDIMP: Percentage of students who did not use an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	93.5 (92.3-94.5)	1,809	88.0 (83.7-91.3)	299	●		
QNOTHHPL: Percentage of students who did not use birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	75.5 (73.5-77.5)	1,809	73.1 (67.7-77.9)	299			●
QNDUALBC: Percentage of students who did not use both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	91.7 (90.3-93.0)	1,786	93.8 (90.3-96.1)	293			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Sexual Behaviors**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy during last sexual intercourse (among students who were currently sexually active)	17.3 (15.5-19.3)	1,809	23.5 (18.8-28.9)	299		●	
QN85: Percentage of students who were not tested for a sexually transmitted disease (STD) other than HIV, such as chlamydia or gonorrhea (during the 12 months before the survey)	62.9 (61.2-64.5)	3,626	62.0 (59.0-65.0)	1,095			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Dietary Behaviors**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>‡</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>‡</sup>	No Difference <sup>‡</sup>
	% 95% CI*	N	% 95% CI	N			
QNOBESE: Percentage of students who had obesity ( $\geq$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) <sup>¶</sup>	15.2 (14.0-16.5)	3,523	20.1 (17.7-22.8)	1,033		●	
QNOWT: Percentage of students who were overweight ( $\geq$ 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) <sup>¶</sup>	17.3 (16.0-18.6)	3,523	18.8 (16.4-21.4)	1,033			●
QN67: Percentage of students who described themselves as slightly or very overweight	24.5 (23.1-25.9)	3,878	33.8 (31.0-36.7)	1,165		●	
QN68: Percentage of students who were not trying to lose weight	61.9 (60.3-63.5)	3,859	50.8 (47.7-53.8)	1,151	●		
QN70: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)	13.9 (12.8-15.1)	3,837	17.3 (15.1-19.7)	1,158		●	

\*Confidence interval.

<sup>‡</sup>Based on t-test analyses,  $p < 0.05$ .

<sup>¶</sup>Based on reference data from the 2000 CDC Growth Charts. In 2017, new, slightly different ranges were used to calculate biologically implausible responses to height and weight questions.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Dietary Behaviors**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN71: Percentage of students who did not eat green salad (one or more times during the 7 days before the survey)	46.2 (44.6-47.9)	3,822	42.2 (39.3-45.2)	1,159	●		
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	17.5 (16.2-18.8)	3,782	18.0 (15.7-20.4)	1,149		●	
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	12.3 (11.2-13.5)	3,782	13.1 (11.1-15.3)	1,149			●
QN77: Percentage of students who did not eat breakfast (during the 7 days before the survey)	17.2 (15.9-18.5)	3,796	22.5 (20.1-25.2)	1,158		●	
QNBK7DAY: Percentage of students who did not eat breakfast on all 7 days (during the 7 days before the survey)	76.6 (75.2-78.0)	3,796	82.7 (80.3-84.8)	1,158		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Physical Activity**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN78: Percentage of students who were not physically active at least 60 minutes per day on 5 or more days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	66.1 (64.5-67.6)	3,747	79.1 (76.6-81.4)	1,145		●	
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	24.3 (22.9-25.7)	3,747	32.4 (29.6-35.3)	1,145		●	
QNPA7DAY: Percentage of students who were not physically active at least 60 minutes per day on all 7 days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	81.1 (79.8-82.4)	3,747	89.7 (87.7-91.4)	1,145		●	
QN79: Percentage of students who watched television 3 or more hours per day (on an average school day)	25.8 (24.4-27.3)	3,714	26.7 (24.1-29.5)	1,136			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.



**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Physical Activity**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN80: Percentage of students who played video or computer games or used a computer 3 or more hours per day (counting time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media, for something that was not school work, on an average school day)	42.0 (40.3-43.7)	3,678	41.4 (38.5-44.4)	1,135			●
QN82: Percentage of students who did not play on at least one sports team (counting any teams run by their school or community groups, during the 12 months before the survey)	43.4 (41.8-45.1)	3,692	48.9 (45.8-51.9)	1,119		●	
QN83: Percentage of students who had a concussion from playing a sport or being physically active (one or more times during the 12 months before the survey)	15.8 (14.6-17.1)	3,649	23.5 (21.0-26.3)	1,120		●	
QN88: Percentage of students who did not get 8 or more hours of sleep (on an average school night)	80.8 (79.4-82.1)	3,645	86.8 (84.7-88.8)	1,130		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

Other	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
Health Risk Behavior							
QN84: Percentage of students who were never tested for human immunodeficiency virus (HIV) (not including tests done when donating blood)	58.2 (56.5-59.8)	3,661	62.6 (59.5-65.5)	1,089		●	
QN86: Percentage of students who did not see a dentist (for a check-up, exam, teeth cleaning, or other dental work, during the 12 months before the survey)	25.2 (23.7-26.7)	3,663	29.4 (26.7-32.3)	1,134		●	
QNNODNT: Percentage of students who never saw a dentist (for a check-up, exam, teeth cleaning, or other dental work)	1.3 (1.0-1.8)	3,663	3.1 (2.1-4.3)	1,134		●	
QN89: Percentage of students who described their grades in school as mostly C's, D's, or F's (during the 12 months before the survey)	38.4 (36.8-40.1)	3,601	36.7 (33.8-39.7)	1,114			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Site-Added**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN93: Percentage of students who are transgender	1.2 (0.9-1.6)	3,907	3.7 (2.7-5.0)	1,178		●	
QN94: Percentage of students who rarely or never wore a helmet when they rode a bicycle or used rollerblades or a skateboard (among students who rode a bicycle or used rollerblades or a skateboard during the 12 months before the survey)	86.2 (84.7-87.5)	2,175	82.5 (79.2-85.3)	600	●		
QN95: Percentage of students who have been afraid of being beaten up at school one or more times (during the 12 months before the survey)	6.7 (5.9-7.5)	3,947	16.6 (14.5-18.9)	1,199		●	
QN96: Percentage of students who reported that their property had been stolen or deliberately damaged on school property one or more times (such as their car, clothing, or books, during the 12 months before the survey)	15.7 (14.5-16.9)	3,929	29.4 (26.8-32.2)	1,196		●	
QN97: Percentage of students who have been harassed on school property one or more times because someone thought they were gay, lesbian, bisexual, or transgender (during the 12 months before the survey)	6.8 (6.0-7.6)	3,932	24.8 (22.3-27.4)	1,199		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Site-Added**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN98: Percentage of students who have been a member of a gang or crew (during the 12 months before the survey)	15.2 (14.1-16.5)	3,814	15.7 (13.7-18.1)	1,108			●
QN99: Percentage of students who have seen or heard people where they live be violent and abusive (including serious hitting, shouting, throwing items, yelling, or name calling, but not 'play fighting,' during the 12 months before the survey)	39.5 (37.9-41.1)	3,896	45.6 (42.6-48.6)	1,141		●	
QN100: Percentage of students who reported someone they were dating or going out with purposely tried to control them or emotionally hurt them one or more times (such things as being told who they could and could not spend time with, being humiliated in front of others, or being threatened if they did not do what they wanted, during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)	18.3 (17.0-19.7)	3,215	36.8 (33.6-40.1)	904		●	
QN101: Percentage of students who reported someone did sexual things to them that they did not want to do one or more times (by pressuring them, lying to them, making promises about the future, threatening to end their relationship, or threatening to spread rumors about them, during the 12 months before the survey)	15.2 (14.1-16.4)	3,934	27.6 (25.1-30.4)	1,177		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Site-Added**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN102: Percentage of students who bullied someone on school property (during the 12 months before the survey)	8.9 (8.0-9.8)	3,892	16.7 (14.6-19.1)	1,128		●	
QN103: Percentage of students who currently used illegal drugs (one or more times during the 30 days before the survey)	14.5 (13.4-15.6)	3,931	28.5 (26.0-31.3)	1,186		●	
QN104: Percentage of students who reported their partners were three or more years older than themselves the last time they had sexual intercourse (among students who have had sexual intercourse)	8.0 (7.1-9.1)	3,090	16.1 (13.6-19.0)	817		●	
QN105: Percentage of students who have had oral sex with one or more people during their life	62.0 (60.4-63.6)	3,905	71.2 (68.5-73.9)	1,185		●	
QN106: Percentage of students who have been pregnant or gotten someone pregnant (one or more times)	7.2 (6.4-8.2)	3,887	10.9 (9.1-13.0)	1,139		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Site-Added**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN108: Percentage of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight (also called fasting, during the 30 days before the survey)	13.3 (12.2-14.5)	3,825	22.6 (20.1-25.3)	1,084		●	
QN109: Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight (during the 30 days before the survey)	5.9 (5.2-6.8)	3,764	13.2 (11.2-15.4)	1,079		●	
QN110: Percentage of students who ate at a fast food chain or carry out restaurant (at least one day during the 7 days before the survey)	82.1 (80.8-83.3)	3,780	82.5 (80.1-84.6)	1,153			●
QN111: Percentage of students who most of the time or always went hungry because there was not enough food in their home (during the 30 days before the survey)	4.2 (3.6-4.9)	3,722	7.2 (5.7-8.9)	1,137		●	
QN112: Percentage of students who reported that none of their classroom teachers provide short physical activity breaks during regular class time (not counting their physical education teacher)	80.1 (78.7-81.5)	3,593	79.2 (76.5-81.7)	1,048			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Site-Added**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN113: Percentage of students who have not been taught or are not sure if they have been taught about AIDS or HIV infection in school	25.6 (24.2-27.1)	3,660	31.2 (28.4-34.2)	1,103		●	
QN114: Percentage of students who strongly disagree, disagree, or are not sure that they feel close to people at their school	34.3 (32.7-35.9)	3,635	39.3 (36.4-42.3)	1,128		●	
QN115: Percentage of students who reported that there is not at least one teacher or other adult in that school that they can talk to if they have a problem or they were not sure if there is at least one teacher or other adult in that school that they can talk to if they have a problem	27.3 (25.8-28.9)	3,576	30.4 (27.6-33.3)	1,082			●
QN116: Percentage of students who have been removed from class for at least one day for disciplinary reasons (counting time spent in in-school suspension, out-of-school suspension, and expulsion, one or more times during the 12 months before the survey)	35.2 (33.6-36.9)	3,641	39.7 (36.7-42.7)	1,122		●	
QN117: Percentage of students who did not usually sleep in their parent's or guardian's home (during the 30 days before the survey)	7.3 (6.5-8.3)	3,629	16.7 (14.5-19.2)	1,126		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

**2019 YOUTH RISK BEHAVIOR SURVEY RESULTS**  
**District of Columbia (Including Charter Schools) High School Survey**  
**Risk Behaviors and Sex of Sexual Contacts Report**

**Site-Added**

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN118: Percentage of students who have ever slept away from their parents or guardians because they were kicked out, ran away, or were abandoned (during the 30 days before the survey)	9.7 (8.7-10.8)	3,563	16.2 (14.0-18.7)	1,053		●	
QN119: Percentage of students who have ever been given money, a place to stay, food, or something else of value in exchange for sex	5.6 (4.8-6.5)	3,549	10.5 (8.7-12.7)	1,037		●	
QN120: Percentage of students who sometimes, rarely, or never get the kind of help they need (among students who report having felt sad, empty, hopeless, angry, or anxious)	75.9 (74.1-77.6)	2,358	78.7 (75.8-81.3)	851			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.