District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

	Heteros	exual	Gay, Lesb Bisext	,			
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN9: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle, one or more times during the 30 days before the survey)	18.7 (17.8-19.6)	8,117	23.6 (21.5-25.8)	1,659		•	
QN10: Percentage of students who drove a car or other vehicle when they had been drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	4.9 (4.1-5.8)	2,861	9.6 (7.3-12.5)	512		•	
QN12: Percentage of students who carried a weapon (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)	15.0 (14.2-15.8)	7,933	17.6 (15.7-19.7)	1,585		•	
QN13: Percentage of students who carried a weapon on school property (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)	4.4 (3.9-4.9)	8,006	6.3 (5.1-7.7)	1,593		•	
QN15: Percentage of students who did not go to school because they felt unsafe at school or on their way to or from school (on at least 1 day during the 30 days before the survey)	7.6 (7.0-8.3)	8,040	12.6 (11.0-14.4)	1,641		•	

^{*}Confidence interval. †Based on t-test analyses, p<0.05.

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

	Heteros	exual	Gay, Lesh Bisext				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN16: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club, one or more times during the 12 months before the survey)	7.7 (7.1-8.4)	8,129	12.6 (11.0-14.3)	1,659		•	
QN17: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)	27.9 (26.8-28.9)	7,849	33.5 (31.0-36.0)	1,577		•	
QN18: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	13.4 (12.6-14.2)	7,956	17.4 (15.5-19.5)	1,600		•	
QN19: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	8.7 (8.0-9.3)	8,000	18.2 (16.3-20.3)	1,583		•	
QN20: Percentage of students who experienced sexual violence (being forced by anyone to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey)	11.5 (10.8-12.3)	7,847	21.4 (19.4-23.6)	1,574		•	

^{*}Confidence interval. †Based on t-test analyses, p<0.05.

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

	Heteros	exual	Gay, Lesh Bisext	,			
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN21: Percentage of students who experienced sexual dating violence (being forced by someone they were dating or going out with to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)	4.2 (3.6-4.8)	4,811	10.1 (8.3-12.1)	1,016		•	
QN22: Percentage of students who experienced physical dating violence (being physically hurt on purpose by someone they were dating or going out with [counting such things as being hit, slammed into something, or injured with an object or weapon] one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)	7.8 (7.1-8.6)	4,944	15.6 (13.5-18.0)	1,072		•	
QN23: Percentage of students who were bullied on school property (ever during the 12 months before the survey)	10.7 (10.1-11.5)	7,871	19.7 (17.8-21.8)	1,576		•	
QN24: Percentage of students who were electronically bullied (counting being bullied through texting, Instagram, Facebook, or other social media, ever during the 12 months before the survey)	8.8 (8.2-9.5)	7,847	16.2 (14.4-18.2)	1,569		•	

^{*}Confidence interval. †Based on t-test analyses, p<0.05.

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

	Heteros	exual	Gay, Lesb Bisext				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN25: Percentage of students who felt sad or hopeless (almost every day for >=2 weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)	29.1 (28.1-30.2)	7,836	51.9 (49.3-54.4)	1,569		•	
QN26: Percentage of students who seriously considered attempting suicide (ever during the 12 months before the survey)	14.7 (14.0-15.6)	7,812	38.7 (36.2-41.2)	1,570		•	
QN27: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)	14.4 (13.6-15.2)	7,837	32.9 (30.6-35.4)	1,574		•	
QN28: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	11.7 (10.9-12.6)	6,499	26.0 (23.7-28.5)	1,372		•	
QN29: Percentage of students who had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	4.6 (4.0-5.1)	6,409	9.6 (8.1-11.4)	1,332		•	

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Tobacco Use

	Heteros	exual	Gay, Lesh Bisext				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN31: Percentage of students who first tried cigarette smoking before age 13 years (even one or two puffs)	8.3 (7.7-9.0)	7,782	13.0 (11.4-14.8)	1,577		•	
QN32: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	4.1 (3.7-4.6)	7,766	7.9 (6.6-9.3)	1,580		•	
QNFRCIG: Percentage of students who currently smoked cigarettes frequently (on 20 or more days during the 30 days before the survey)	0.5 (0.3-0.7)	7,766	1.3 (0.8-2.0)	1,580		•	
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	0.3 (0.2-0.5)	7,766	1.0 (0.6-1.6)	1,580		•	
QN35: Percentage of students who currently used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo], on at least 1 day during the 30 days before the survey)	11.3 (10.6-12.1)	7,746	17.1 (15.3-19.0)	1,586		•	

^{*}Confidence interval. †Based on t-test analyses, p<0.05.

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Tobacco Use

	Hetero	sexual	Gay, Lesh Bisex				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QNFREVP: Percentage of students who currently used electronic vapor products frequently (on 20 or more days during the 30 days before the survey)	1.4 (1.1-1.6)	7,746	1.8 (1.2-2.5)	1,586			•
QNDAYEVP: Percentage of students who currently used electronic vapor products daily (on all 30 days during the 30 days before the survey)	0.8 (0.6-1.0)	7,746	1.2 (0.8-1.9)	1,586			•
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Copenhagen, Grizzly, Skoal, or Camel Snus], not counting any electronic vapor products, on at least 1 day during the 30 days before the survey)	4.0 (3.6-4.5)	7,844	8.1 (6.8-9.6)	1,612		•	
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars, on at least 1 day during the 30 days before the survey)	5.3 (4.8-5.8)	7,815	8.3 (7.1-9.8)	1,610		•	
QNTB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)	7.0 (6.5-7.7)	7,758	12.1 (10.6-13.8)	1,598		•	

^{*}Confidence interval. †Based on t-test analyses, p<0.05.

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Tobacco Use

	Heteros	exual	Gay, Lesh Bisext				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QNTB3: Percentage of students who currently smoked cigarettes or cigars or used smokeless tobacco (on at least 1 day during the 30 days before the survey)	8.2 (7.5-8.8)	7,778	14.4 (12.7-16.2)	1,603		•	
QNTB4: Percentage of students who currently smoked cigarettes or cigars or used smokeless tobacco or electronic vapor products (on at least 1 day during the 30 days before the survey)	14.9 (14.1-15.7)	7,705	22.7 (20.7-24.9)	1,592		•	
QNTB5: Percentage of students who currently smoked cigarettes or used electronic vapor products (on at least 1 day during the 30 days before the survey)	12.9 (12.2-13.7)	7,689	20.0 (18.0-22.1)	1,578		•	

^{*}Confidence interval.
*Based on t-test analyses, p<0.05.

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

	Heteros	exual	Gay, Lesh Bisex				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN40: Percentage of students who had their first drink of alcohol before age 13 years (other than a few sips)	16.3 (15.4-17.2)	7,484	23.4 (21.2-25.7)	1,527		•	
QN41: Percentage of students who currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)	17.8 (16.9-18.7)	7,427	30.8 (28.4-33.2)	1,493		•	
QN42: Percentage of students who currently were binge drinking (had four or more drinks of alcohol in a row for female students or five or more drinks of alcohol in a row for male students, within a couple of hours, on at least 1 day during the 30 days before the survey)	7.1 (6.5-7.7)	7,593	13.0 (11.5-14.8)	1,534		•	
QN46: Percentage of students who tried marijuana for the first time before age 13 years	10.7 (10.0-11.5)	7,592	14.6 (12.8-16.5)	1,554		•	
QN47: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	27.1 (26.1-28.1)	7,656	40.6 (38.1-43.2)	1,573		•	

^{*}Confidence interval. †Based on t-test analyses, p<0.05.

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

	Heteros	exual	Gay, Lesb Bisext				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN49: Percentage of students who ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)	17.5 (16.6-18.4)	7,728	28.3 (26.0-30.6)	1,595		•	
QN50: Percentage of students who ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)	4.4 (3.9-4.9)	7,653	8.6 (7.3-10.1)	1,582		•	
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	8.9 (8.3-9.6)	7,720	15.9 (14.2-17.9)	1,587		•	
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)	3.7 (3.3-4.2)	7,700	7.2 (6.0-8.7)	1,586		•	
QN53: Percentage of students who ever used methamphetamines (also called "speed," "crystal meth," "crank," "ice," or "meth," one or more times during their life)	3.8 (3.4-4.3)	7,644	7.1 (5.8-8.5)	1,572		•	

^{*}Confidence interval. †Based on t-test analyses, p<0.05.

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

	Heteros	sexual	Gay, Lesl Bisex				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)	4.2 (3.7-4.7)	7,653	7.1 (6.0-8.5)	1,580		•	

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

	Heteros	exual	Gay, Lesb Bisext				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN58: Percentage of students who ever had sexual intercourse	43.8 (42.6-45.1)	6,478	48.5 (45.6-51.3)	1,302		•	
QN59: Percentage of students who had sexual intercourse for the first time before age 13 years	8.3 (7.6-9.1)	6,509	5.6 (4.5-7.1)	1,316	•		
QN60: Percentage of students who had sexual intercourse with four or more persons during their life	12.4 (11.5-13.3)	6,447	11.9 (10.2-13.8)	1,305			•
QN61: Percentage of students who were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)	30.3 (29.2-31.5)	6,416	35.1 (32.4-37.8)	1,302		•	
QN62: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	15.4 (13.8-17.2)	1,820	19.7 (16.3-23.7)	440		•	

^{*}Confidence interval. †Based on t-test analyses, p<0.05.

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

	Heteros	exual	Gay, Lesb Bisext				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN63: Percentage of students who did not use a condom during last sexual intercourse (among students who were currently sexually active)	39.8 (37.4-42.2)	1,764	57.1 (52.2-61.9)	427		•	
QN64: Percentage of students who did not use birth control pills before last sexual intercourse (to prevent pregnancy among students who were currently sexually active)	90.1 (88.6-91.4)	1,760	91.7 (88.7-93.9)	428			•
QNIUDIMP: Percentage of students who did not use an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	93.2 (92.0-94.3)	1,760	91.3 (88.2-93.6)	428			•
QNOTHHPL: Percentage of students who did not use birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	76.0 (73.9-77.9)	1,760	75.8 (71.4-79.7)	428			•
QNDUALBC: Percentage of students who did not use both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	92.3 (90.9-93.5)	1,727	92.8 (89.7-95.1)	414			•

^{*}Confidence interval. †Based on t-test analyses, p<0.05.

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

	Heteros	exual	Gay, Lesbian, or Bisexual				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy during last sexual intercourse (among students who were currently sexually active)	17.5 (15.7-19.5)	1,760	35.5 (30.9-40.4)	428		•	
QN85: Percentage of students who were not tested for a sexually transmitted disease (STD) other than HIV, such as chlamydia or gonorrhea (during the 12 months before the survey)	72.6 (71.5-73.7)	6,962	69.3 (66.7-71.7)	1,450	•		

District of Columbia (Including Charter Schools) High School Survey **Risk Behaviors and Sexual Identity Report**

Dietary Behaviors

	Heterosexual Gay, Lesbian, or Bisexual		,				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QNOBESE: Percentage of students who had obesity (>= 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [¶]	16.1 (15.2-17.0)	7,064	20.5 (18.3-22.9)	1,410		•	
QNOWT: Percentage of students who were overweight (>= 85th percentile but <95th percentile for body mass index, based on sexand age-specific reference data from the 2000 CDC growth charts)¶	17.1 (16.2-18.0)	7,064	20.8 (18.7-23.1)	1,410		•	
QN67: Percentage of students who described themselves as slightly or very overweight	24.9 (23.9-26.0)	7,379	34.8 (32.4-37.3)	1,525		•	
QN68: Percentage of students who were not trying to lose weight	58.5 (57.4-59.7)	7,354	46.5 (43.9-49.1)	1,515	•		
QN70: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)	15.5 (14.6-16.4)	7,348	16.7 (14.7-18.8)	1,523			•

^{*}Confidence interval. *Based on t-test analyses, p<0.05.

Based on reference data from the 2000 CDC Growth Charts. In 2017, new, slightly different ranges were used to calculate biologically implausible responses to height and weight questions.

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

	Heterosexual		Gay, Lesbian, or Bisexual				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN71: Percentage of students who did not eat green salad (one or more times during the 7 days before the survey)	46.6 (45.4-47.8)	7,321	45.9 (43.3-48.5)	1,525			•
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	15.2 (14.4-16.1)	7,255	14.2 (12.4-16.1)	1,504			•
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	10.4 (9.7-11.2)	7,255	10.0 (8.5-11.8)	1,504			•
QN77: Percentage of students who did not eat breakfast (during the 7 days before the survey)	17.9 (17.0-18.9)	7,282	20.8 (18.7-23.1)	1,523		•	
QNBK7DAY: Percentage of students who did not eat breakfast on all 7 days (during the 7 days before the survey)	73.6 (72.5-74.6)	7,282	81.0 (78.9-82.9)	1,523		•	

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Physical Activity

	Heteros	exual	Gay, Lesb Bisext				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN78: Percentage of students who were not physically active at least 60 minutes per day on 5 or more days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	70.6 (69.5-71.7)	7,215	77.3 (75.0-79.4)	1,502		•	
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	27.0 (25.9-28.1)	7,215	32.0 (29.6-34.6)	1,502		•	
QNPA7DAY: Perentage of students who were not physically active at least 60 minutes per day on all 7 days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	84.2 (83.3-85.1)	7,215	88.6 (86.8-90.2)	1,502		•	
QN79: Percentage of students who watched television 3 or more hours per day (on an average school day)	24.9 (23.8-26.0)	7,167	25.4 (23.1-27.8)	1,493			•

^{*}Confidence interval. †Based on t-test analyses, p<0.05.

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Physical Activity

	Heteros	exual	Gay, Lesb Bisext	,			
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN80: Percentage of students who played video or computer games or used a computer 3 or more hours per day (counting time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media, for something that was not school work, on an average school day)	41.0 (39.8-42.2)	7,108	42.8 (40.2-45.5)	1,489			•
QN82: Percentage of students who did not play on at least one sports team (counting any teams run by their school or community groups, during the 12 months before the survey)	48.6 (47.4-49.8)	7,101	53.1 (50.4-55.7)	1,475		•	
QN83: Percentage of students who had a concussion from playing a sport or being physically active (one or more times during the 12 months before the survey)	14.6 (13.8-15.5)	7,014	15.7 (13.9-17.8)	1,464			•
QN88: Percentage of students who did not get 8 or more hours of sleep (on an average school night)	78.2 (77.1-79.1)	7,011	83.2 (81.1-85.1)	1,485		•	

^{*}Confidence interval. †Based on t-test analyses, p<0.05.

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Other

	Heterosexual		Gay, Lesbian, or Bisexual				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN84: Percentage of students who were never tested for human immunodeficiency virus (HIV) (not including tests done when donating blood)	68.5 (67.3-69.6)	7,026	68.0 (65.4-70.4)	1,454			•
QN86: Percentage of students who did not see a dentist (for a check-up, exam, teeth cleaning, or other dental work, during the 12 months before the survey)	24.6 (23.5-25.7)	7,068	26.3 (24.0-28.7)	1,482			•
QNNODNT: Percentage of students who never saw a dentist (for a check-up, exam, teeth cleaning, or other dental work)	1.7 (1.4-2.1)	7,068	2.4 (1.7-3.4)	1,482			•
QN89: Percentage of students who described their grades in school as mostly C's, D's, or F's (during the 12 months before the survey)	33.8 (32.6-35.0)	6,950	34.5 (32.0-37.1)	1,462			•

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

	Heteros	Gay, Lesbian, or sexual Bisexual					
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN93: Percentage of students who are transgender	1.0 (0.8-1.3)	8,082	4.0 (3.1-5.0)	1,644		•	
QN94: Percentage of students who rarely or never wore a helmet when they rode a bicycle or used rollerblades or a skateboard (among students who rode a bicycle or used rollerblades or a skateboard during the 12 months before the survey)	82.6 (81.5-83.7)	4,314	77.7 (74.7-80.5)	758	•		
QN95: Percentage of students who have been afraid of being beaten up at school one or more times (during the 12 months before the survey)	7.1 (6.6-7.7)	8,141	12.8 (11.2-14.5)	1,663		•	
QN96: Percentage of students who reported that their property had been stolen or deliberately damaged on school property one or more times (such as their car, clothing, or books, during the 12 months before the survey)	15.0 (14.2-15.9)	8,119	21.9 (19.9-24.0)	1,655		•	
QN97: Percentage of students who have been harassed on school property one or more times because someone thought they were gay, lesbian, bisexual, or transgender (during the 12 months before the survey)	6.4 (5.9-7.0)	8,125	20.6 (18.6-22.7)	1,660		•	

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

	Heteros	exual	Gay, Lesh Bisex				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN98: Percentage of students who have been a member of a gang or crew (during the 12 months before the survey)	13.3 (12.5-14.1)	7,878	13.1 (11.5-14.9)	1,572			•
QN99: Percentage of students who have seen or heard people where they live be violent and abusive (including serious hitting, shouting, throwing items, yelling, or name calling, but not 'play fighting,' during the 12 months before the survey)	35.0 (33.9-36.1)	8,023	43.4 (40.9-45.9)	1,610		•	
QN100: Percentage of students who reported someone they were dating or going out with purposely tried to control them or emotionally hurt them one or more times (such things as being told who they could and could not spend time with, being humiliated in front of others, or being threatened if they did not do what they wanted, during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)	16.0 (15.0-17.1)	5,205	32.6 (29.8-35.5)	1,156		•	
QN101: Percentage of students who reported someone did sexual things to them that they did not want to do one or more times (by pressuring them, lying to them, making promises about the future, threatening to end their relationship, or threatening to spread rumors about them, during the 12 months before the survey)	12.0 (11.3-12.8)	8,015	20.3 (18.4-22.4)	1,632		•	

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

	· · · · · · · · · · · · · · · · · ·		Gay, Lesh Bisext				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN102: Percentage of students who bullied someone on school property (during the 12 months before the survey)	8.7 (8.1-9.4)	7,889	12.6 (11.1-14.4)	1,586		•	
QN103: Percentage of students who currently used illegal drugs (one or more times during the 30 days before the survey)	11.1 (10.4-11.8)	7,636	18.7 (16.8-20.7)	1,577		•	
QN104: Percentage of students who reported their partners were three or more years older than themselves the last time they had sexual intercourse (among students who have had sexual intercourse)	8.4 (7.5-9.5)	3,181	13.9 (11.6-16.7)	761		•	
QN105: Percentage of students who have had oral sex with one or more people during their life	35.8 (34.7-37.0)	7,416	49.7 (47.1-52.3)	1,545		•	
QN106: Percentage of students who have been pregnant or gotten someone pregnant (one or more times)	5.0 (4.5-5.5)	7,353	6.2 (5.0-7.6)	1,517			•

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

	Heteros	exual	Gay, Lesh Bisext				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN108: Percentage of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight (also called fasting, during the 30 days before the survey)	13.6 (12.8-14.4)	7,288	21.5 (19.4-23.8)	1,475		•	
QN109: Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight (during the 30 days before the survey)	6.5 (5.9-7.1)	7,189	9.8 (8.4-11.6)	1,450		•	
QN110: Percentage of students who ate at a fast food chain or carry out restaurant (at least one day during the 7 days before the survey)	78.3 (77.3-79.3)	7,247	80.6 (78.5-82.6)	1,517			•
QN111: Percentage of students who most of the time or always went hungry because there was not enough food in their home (during the 30 days before the survey)	4.2 (3.7-4.7)	7,158	4.8 (3.8-6.1)	1,487			•
QN112: Percentage of students who reported that none of their classroom teachers provide short physical activity breaks during regular class time (not counting their physical education teacher)	79.4 (78.4-80.4)	6,928	81.4 (79.2-83.4)	1,420			•

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

	Heteros	exual	Gay, Lesbian, or Bisexual				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN113: Percentage of students who have not been taught or are not sure if they have been taught about AIDS or HIV infection in school	28.3 (27.2-29.4)	7,042	29.2 (26.8-31.7)	1,468			•
QN114: Percentage of students who strongly disagree, disagree, or are not sure that they feel close to people at their school	33.2 (32.1-34.4)	7,029	38.3 (35.7-40.9)	1,480		•	
QN115: Percentage of students who reported that there is not at least one teacher or other adult in that school that they can talk to if they have a problem or they were not sure if there is at least one teacher or other adult in that school that they can talk to if they have a problem	28.3 (27.2-29.4)	6,923	25.9 (23.6-28.3)	1,442			•
QN116: Percentage of students who have been removed from class for at least one day for disciplinary reasons (counting time spent in in-school suspension, out-of-school suspension, and expulsion, one or more times during the 12 months before the survey)	28.6 (27.5-29.8)	7,020	33.3 (30.8-35.9)	1,475		•	
QN117: Percentage of students who did not usually sleep in their parent's or guardian's home (during the 30 days before the survey)	7.3 (6.7-8.0)	7,005	10.5 (9.0-12.3)	1,476		•	

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

	Heteros	exual	Gay, Lesh Bisex				
Health Risk Behavior	% 95% CI*	N	% 95% CI	N	Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
QN118: Percentage of students who have ever slept away from their parents or guardians because they were kicked out, ran away, or were abandoned (during the 30 days before the survey)	9.7 (9.0-10.5)	6,861	11.8 (10.1-13.7)	1,422		•	
QN119: Percentage of students who have ever been given money, a place to stay, food, or something else of value in exchange for sex	6.4 (5.8-7.1)	6,841	6.9 (5.6-8.4)	1,414			•
QN120: Percentage of students who sometimes, rarely, or never get the kind of help they need (among students who report having felt sad, empty, hopeless, angry, or anxious)	74.3 (72.9-75.6)	4,227	75.5 (72.8-77.9)	1,166			•