

CHILD CARE: CORONAVIRUS (COVID-19) ISOLATION AND RETURN REQUIREMENTS

Last Updated: July 15, 2022

ISOLATION: Physically separating an individual with symptoms or confirmed diagnosis of COVID-19 away from others to prevent spread. COVID-19 typically spreads between individuals who are in close contact (within about 6 feet) through respiratory droplets created when someone talks, coughs, or sneezes.

If your child has tested positive for COVID-19 or has symptoms of COVID-19, immediately report it to your child care facility and begin isolation. The length of isolation will depend on an individual's age, severity of symptoms and whether the positive individual can correctly and consistently wear a well-fitting mask.

This resource applies to any individual regardless of vaccination status or history of COVID-19 diagnosis. District of Columbia regulations for child development facilities require children and staff to be temporarily removed from care if they have symptoms or have tested positive for a communicable disease.

WHAT TO DO WHILE IN ISOLATION:

- Stay at home (if it is safe to do so), except to get medical care.
- Do not travel.
- Separate yourself from other people and pets in your home as much as possible, and if possible, in a separate room and bathroom (or clean between uses).
 - Especially separate from anyone who is immunocompromised, at higher risk for severe COVID-19, or unvaccinated or not up to date on their COVID-19 vaccination.
- Wear a well-fitting mask if you need to be near any other members of your household (caregivers should also wear a mask if they need to be near the isolated person. Note: masks should not be worn while sleeping or on children under age 2).
- Bring as much fresh air into your home as possible, weather and safety permitting, especially in any shared spaces.
- Eat in separate areas, and wash dishes used by the sick person in hot water while wearing gloves or in dishwasher.
- Do not share personal items (dishes, towels, cell phones, etc.) with others.
- Do not allow visitors in your home.
- Cover your coughs and sneezes.
- Wash your hands frequently.
- Monitor symptoms and seek emergency medical attention if the ill individual has any of the following symptoms:
 - trouble breathing
 - not enough breath to get out a full sentence
 - coughing so much they can't catch their breath
 - persistent pain or pressure in the chest
 - new confusion
 - inability to wake/stay awake
 - pale/gray/blue-colored lips or nail beds (depending on skin tone)
 - *This list does not include all possible symptoms. Please call your medical provider for any other symptoms that are severe or concerning to you. Call 911 or call ahead to your local emergency facility if necessary and notify the operator that you are seeking care for someone who has or may have COVID-19.*



CALCULATING ISOLATION: Day 0 is the first day of symptoms or a positive viral test. Day 1 is the first full day after symptoms developed or the specimen was collected.

WHEN SHOULD MY CHILD RETURN TO CHILD CARE IF I AM UNSURE WHETHER THE SYMPTOMS ARE DUE TO COVID-19?

RETURNING AFTER SYMPTOMS OF COVID-19	
IF TESTED FOR COVID-19	IF NOT TESTED FOR COVID-19
<ul style="list-style-type: none"> If test is positive: See table below. If test is negative: When the child care facility's standard criteria to return after illness is met. Note: You may use either an antigen test or a NAAT (e.g., PCR) test. 	<ul style="list-style-type: none"> Submit documentation from a healthcare provider of an alternate diagnosis and meet standard criteria to return after illness. <p>OR</p> <ul style="list-style-type: none"> Meet COVID-19 symptom-based criteria to return: <ul style="list-style-type: none"> At least 24 hours after the fever has resolved without the use of fever-reducing medication (e.g., acetaminophen, ibuprofen) and symptoms have improved; <p>AND</p> <ul style="list-style-type: none"> Under age 2 <u>or</u> unable to wear a well-fitting mask: After at least 10 days from when symptoms first started (i.e., on day 11) Age 2 or older <u>and</u> able to wear a well-fitting mask: After at least 5 days from when symptoms first started (i.e., on day 6) and able to wear a well-fitting mask while in child care through day 10

WHEN SHOULD MY CHILD RETURN TO CHILD CARE AFTER TESTING POSITIVE?

RETURNING FROM ISOLATION AFTER TESTING POSITIVE FOR COVID-19		
SYMPTOM SEVERITY	UNDER AGE 2 OR NOT ABLE TO WEAR A WELL-FITTING MASK	AGE 2 OR OLDER <u>AND</u> ABLE TO WEAR A WELL-FITTING MASK
<p>Mild to Moderate Symptoms of COVID-19</p> <p>Mild to Moderate Symptoms Include:</p> <ul style="list-style-type: none"> Able to recover at home taking over-the-counter medications or oral prescription medications. Did not need major medical intervention during their illness. Did not require emergency room visits or admission to a hospital. Did not develop severe breathing trouble. 	<ul style="list-style-type: none"> Return after at least 10 days (i.e., on day 11) from when symptoms first appeared. <p><i>Note: Children under age 2 have a longer isolation period because they cannot safely wear a mask.</i></p>	<ul style="list-style-type: none"> Return after at least 5 days (i.e., on day 6) from when symptoms first appeared if fever-free for at least 24 hours without the use of fever-lowering medications (e.g., acetaminophen or ibuprofen) AND other symptoms are improving. Wear a well-fitting mask while in child care through day 10. If still have a fever and/or other symptoms have not improved at day 5: <ul style="list-style-type: none"> Return after being fever free for 24 hours without the use of fever-lowering medication and other symptoms have improved. <p>OR</p> <ul style="list-style-type: none"> Return after at least 10 days (i.e., on day 11) from when symptoms first appeared. <p>If an individual has access to a test and wants to re-test, the best approach is to use an antigen test towards the end of the 5-day isolation period. Test only if fever-free for 24 hours without the use of fever-lowering medication and other symptoms have improved.</p> <ul style="list-style-type: none"> Positive test: Remain in isolation and return at least 10 days (i.e., on day 11) from when symptoms first appeared. Negative test: End isolation but continue to wear a well-fitting mask in child care through
<p>Moderate to Severe Symptoms of COVID-19 OR Individual has Weakened Immune System</p> <p>Severe or Critical Symptoms Include:</p> <ul style="list-style-type: none"> Became sicker than the description of mild to moderate COVID-19. Had dangerously low oxygen, or serious trouble breathing. Needed more intensive medical attention such as a trip to the emergency room, admission to the hospital, admission to an intensive care unit, or required mechanical ventilation or "life support" for any amount of time. 	<ul style="list-style-type: none"> Return after at least 10 days (i.e., on day 11) from when symptoms first appeared. <p>Recommended to consult with healthcare provider before returning.</p>	<ul style="list-style-type: none"> Return after at least 10 days (i.e., on day 11) from when symptoms first appeared. <p>Recommended to consult with healthcare provider before returning.</p>
<p>No Symptoms (Asymptomatic Infection)</p>	<ul style="list-style-type: none"> Return after at least 10 days (i.e., on day 11) from date of positive test. <p><i>Note: Children under age 2 have a longer isolation period because they cannot safely wear a mask.</i></p>	<ul style="list-style-type: none"> If continue to have no symptoms: Return after at least 5 days (i.e., on day 6) from positive test and wear a well-fitting mask in child care through day 10 If symptoms develop: Isolation restarts with day 0 as the day symptoms first appeared. <p>If symptoms have not developed and an individual has access to a test and wants to re-test, the best approach is to use an antigen test towards the end of the 5-day isolation period.</p> <ul style="list-style-type: none"> Positive test: Remain in isolation and return at least 10 days (i.e., on day 11) from the initial positive test. Negative test: End isolation but continue to wear a well-fitting mask in child care through day 10.