

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN9: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle, one or more times during the 30 days before the survey)	23.0 (21.4-24.8)	3,000	31.4 (28.1-35.0)	894		●	
QN10: Percentage of students who drove a car or other vehicle when they had been drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	4.7 (3.5-6.3)	1,146	19.1 (14.9-24.2)	343		●	
QN12: Percentage of students who carried a weapon (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)	23.0 (21.4-24.7)	3,024	28.7 (25.6-32.1)	903		●	
QN15: Percentage of students who did not go to school because they felt unsafe at school or on their way to or from school (on at least 1 day during the 30 days before the survey)	7.9 (6.9-9.1)	2,996	15.7 (13.3-18.5)	889		●	
QN16: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club, one or more times during the 12 months before the survey)	8.2 (7.2-9.3)	3,034	18.4 (15.7-21.4)	910		●	
QN17: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)	34.5 (32.6-36.4)	2,963	42.9 (39.3-46.7)	855		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN18: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	16.7 (15.2-18.2)	2,966	21.7 (18.6-25.2)	849		●	
QN19: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	7.7 (6.7-8.8)	3,012	18.7 (16.1-21.7)	863		●	
QN22: Percentage of students who experienced physical dating violence (being physically hurt on purpose by someone they were dating or going out with [counting such things as being hit, slammed into something, or injured with an object or weapon] one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)	12.6 (11.3-14.1)	2,389	24.4 (20.9-28.2)	686		●	
QN23: Percentage of students who were bullied on school property (ever during the 12 months before the survey)	9.4 (8.4-10.6)	3,011	16.8 (14.1-19.8)	861		●	
QN24: Percentage of students who were electronically bullied (counting being bullied through texting, Instagram, Facebook, or other social media, ever during the 12 months before the survey)	7.6 (6.6-8.6)	2,991	14.9 (12.4-17.7)	859		●	
QN25: Percentage of students who felt sad or hopeless (almost every day for >=2 weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)	26.7 (25.0-28.4)	2,990	43.1 (39.5-46.8)	855		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [‡]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [‡]	No Difference [‡]
	% 95% CI [*]	N	% 95% CI	N			
QN26: Percentage of students who seriously considered attempting suicide (ever during the 12 months before the survey)	13.9 (12.6-15.3)	2,988	31.4 (28.1-34.9)	857		●	
QN27: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)	13.6 (12.2-15.0)	2,991	29.2 (25.9-32.7)	857		●	
QN28: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	14.1 (12.6-15.7)	2,504	30.6 (27.0-34.4)	739		●	
QN29: Percentage of students who had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	6.7 (5.7-7.9)	2,483	10.9 (8.5-13.9)	714		●	

^{*}Confidence interval.

[‡]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Tobacco Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN31: Percentage of students who first tried cigarette smoking before age 13 years (even one or two puffs)	13.0 (11.6-14.4)	2,979	21.5 (18.6-24.8)	872		●	
QN34: Percentage of students who ever used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo])	34.1 (32.2-35.9)	2,934	48.3 (44.5-52.1)	825		●	
QN35: Percentage of students who currently used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo], on at least 1 day during the 30 days before the survey)	11.9 (10.7-13.3)	2,983	21.3 (18.4-24.5)	877		●	
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars, on at least 1 day during the 30 days before the survey)	9.9 (8.8-11.2)	3,008	18.9 (16.3-21.7)	900		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Alcohol/Other Drug Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN40: Percentage of students who ever drank alcohol (at least one drink of alcohol, on at least 1 day during their life)	59.2 (57.1-61.2)	2,685	76.4 (72.9-79.5)	758		●	
QN41: Percentage of students who had their first drink of alcohol before age 13 years (other than a few sips)	17.5 (16.0-19.1)	2,683	29.8 (26.2-33.7)	771		●	
QN42: Percentage of students who currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)	26.0 (24.3-27.8)	2,784	40.1 (36.3-44.0)	775		●	
QN44: Percentage of students who currently were binge drinking (had four or more drinks of alcohol in a row for female students or five or more drinks of alcohol in a row for male students, within a couple of hours, on at least 1 day during the 30 days before the survey)	10.2 (9.1-11.4)	2,913	17.7 (14.9-20.8)	824		●	
QN47: Percentage of students who tried marijuana for the first time before age 13 years	17.9 (16.4-19.5)	2,966	28.4 (25.2-31.8)	881		●	
QN48: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	43.1 (41.2-45.1)	2,943	53.9 (50.2-57.6)	873		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Alcohol/Other Drug Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN49: Percentage of students who ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)	5.2 (4.4-6.1)	3,023	16.4 (13.9-19.3)	900		●	
QN50: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	10.5 (9.3-11.8)	2,973	24.8 (21.7-28.2)	881		●	
QN51: Percentage of students who ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)	2.9 (2.3-3.6)	3,025	11.4 (9.3-13.9)	905		●	
QN52: Percentage of students who ever used methamphetamines (also called "speed," "crystal," "crank," or "ice," one or more times during their life)	3.9 (3.2-4.8)	3,015	13.6 (11.2-16.3)	901		●	
QN53: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)	5.4 (4.6-6.4)	2,998	16.6 (14.0-19.6)	896		●	
QN54: Percentage of students who ever used synthetic marijuana (also called "K2," "Spice," "fake weed," "King Kong," "Yucatan Fire," "Skunk," or "Moon Rocks," one or more times during their life)	5.7 (4.8-6.7)	3,021	15.0 (12.6-17.8)	902		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Alcohol/Other Drug Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI [*]	N	% 95% CI	N			
QN56: Percentage of students who ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their	16.2 (14.8-17.8)	2,995	31.4 (28.0-34.9)	894		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Sexual Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN59: Percentage of students who ever had sexual intercourse	84.3 (82.9-85.6)	2,632	73.4 (69.6-76.9)	679	●		
QN60: Percentage of students who had sexual intercourse for the first time before age 13 years	15.5 (13.9-17.2)	2,600	15.9 (13.0-19.4)	687			●
QN61: Percentage of students who had sexual intercourse with four or more persons during their life	25.8 (23.9-27.7)	2,598	22.6 (19.2-26.4)	683			●
QN62: Percentage of students who were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)	57.6 (55.5-59.6)	2,605	51.4 (47.3-55.6)	685	●		
QN63: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	17.0 (14.9-19.3)	1,397	28.5 (23.5-33.9)	337		●	
QN64: Percentage of students who did not use a condom during last sexual intercourse (among students who were currently sexually active)	35.4 (32.7-38.3)	1,358	55.4 (49.3-61.3)	318		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sex of Sexual Contacts Report

Sexual Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN65: Percentage of students who did not use birth control pills before last sexual intercourse (to prevent pregnancy among students who were currently sexually active)	92.0 (90.3-93.4)	1,360	88.9 (84.2-92.3)	328			●
QNIUDIMP: Percentage of students who did not use an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	93.7 (92.2-95.0)	1,360	95.1 (92.4-96.9)	328			●
QNSHPARG: Percentage of students who did not use a shot (e.g., Depo-Provera), patch (e.g., OrthoEvra), or birth control ring (e.g., NuvaRing) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	94.1 (92.6-95.3)	1,360	94.2 (90.6-96.5)	328			●
QNOTHHPL: Percentage of students who did not use birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	79.8 (77.4-82.0)	1,360	78.2 (72.8-82.8)	328			●
QNDUALBC: Percentage of students who did not use both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	92.5 (90.7-94.0)	1,342	91.1 (86.9-94.1)	316			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Sexual Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [‡]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [‡]	No Difference [‡]
	% 95% CI [*]	N	% 95% CI	N			
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy during last sexual intercourse (among students who were currently sexually active)	18.6 (16.4-21.1)	1,360	36.9 (31.4-42.7)	328		●	

^{*}Confidence interval.

[‡]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [‡]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [‡]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QNOBES: Percentage of students who had obesity (\geq 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [‡]	15.2 (13.8-16.8)	2,833	17.6 (14.9-20.7)	818			●
QNOWT: Percentage of students who were overweight (\geq 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [‡]	16.8 (15.3-18.4)	2,833	19.2 (16.3-22.6)	818			●
QN68: Percentage of students who described themselves as slightly or very overweight	22.8 (21.2-24.4)	2,946	30.3 (27.0-33.8)	881		●	
QN69: Percentage of students who were not trying to lose weight	61.8 (59.9-63.7)	2,940	52.2 (48.4-55.9)	866	●		
QN70: Percentage of students who did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)	24.0 (22.4-25.7)	2,858	26.0 (23.0-29.3)	860			●
QN71: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)	15.0 (13.6-16.5)	2,844	14.6 (12.2-17.4)	859			●

*Confidence interval.

[‡]Based on t-test analyses, $p < 0.05$.

[‡]Based on reference data from the 2000 CDC Growth Charts. In 2017, new, slightly different ranges were used to calculate biologically implausible responses to height and weight questions.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNFR0: Percentage of students who did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	6.9 (5.9-8.0)	2,835	7.2 (5.6-9.3)	854			●
QN72: Percentage of students who did not eat green salad (one or more times during the 7 days before the survey)	46.1 (44.1-48.1)	2,812	37.6 (34.0-41.3)	853	●		
QN73: Percentage of students who did not eat potatoes (one or more times during the 7 days before the survey)	48.8 (46.8-50.8)	2,803	46.0 (42.3-49.7)	854			●
QN74: Percentage of students who did not eat carrots (one or more times during the 7 days before the survey)	66.1 (64.2-68.0)	2,791	61.8 (58.1-65.3)	848	●		
QN75: Percentage of students who did not eat other vegetables (one or more times during the 7 days before the survey)	24.3 (22.5-26.1)	2,770	24.6 (21.5-28.1)	837			●
QNVEG0: Percentage of students who did not eat vegetables (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	10.8 (9.6-12.2)	2,725	8.6 (6.7-10.9)	821			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	19.4 (17.7-21.1)	2,782	19.2 (16.3-22.5)	843			●
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	13.9 (12.4-15.5)	2,782	13.9 (11.3-17.0)	843			●
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	8.3 (7.2-9.7)	2,782	7.8 (6.0-10.2)	843			●
QN78: Percentage of students who did not eat breakfast (during the 7 days before the survey)	16.6 (15.1-18.2)	2,738	20.7 (17.8-24.0)	837		●	
QNBK7DAY: Percentage of students who did not eat breakfast on all 7 days (during the 7 days before the survey)	76.6 (74.8-78.3)	2,738	85.5 (82.8-87.8)	837		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Physical Activity

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN79: Percentage of students who were not physically active at least 60 minutes per day on 5 or more days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	69.9 (68.0-71.8)	2,716	80.2 (77.2-82.9)	833		●	
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	25.3 (23.5-27.2)	2,716	32.0 (28.5-35.7)	833		●	
QNPA7DAY: Percentage of students who were not physically active at least 60 minutes per day on all 7 days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	83.5 (81.9-85.0)	2,716	88.5 (85.9-90.6)	833		●	
QN80: Percentage of students who watched television 3 or more hours per day (on an average school day)	30.5 (28.5-32.5)	2,674	27.3 (23.9-30.9)	818			●
QN81: Percentage of students who played video or computer games or used a computer 3 or more hours per day (counting time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media, for something that was not school work, on an average school day)	42.9 (40.9-45.0)	2,719	39.0 (35.4-42.7)	824			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Physical Activity

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI [*]	N	% 95% CI	N			
QN83: Percentage of students who did not play on at least one sports team (counting any teams run by their school or community groups, during the 12 months before the survey)	41.4 (39.4-43.5)	2,702	48.3 (44.5-52.1)	814		●	
QN84: Percentage of students who had a concussion from playing a sport or being physically active (one or more times during the 12 months before the survey)	17.4 (15.9-19.0)	2,713	22.8 (19.8-26.1)	817		●	
QN88: Percentage of students who did not get 8 or more hours of sleep (on an average school night)	78.0 (76.4-79.6)	3,032	80.9 (77.8-83.6)	908			●

^{*}Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Other	Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
		% 95% CI [*]	N	% 95% CI	N			
	QN85: Percentage of students who were never tested for human immunodeficiency virus (HIV) (not including tests done when donating blood)	62.3 (60.3-64.2)	2,692	59.1 (55.3-62.8)	807			●
	QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma	33.6 (31.7-35.6)	2,691	39.3 (35.6-43.2)	800		●	
	QN89: Percentage of students who described their grades in school as mostly C's, D's, or F's (during the 12 months before the survey)	43.7 (41.8-45.7)	3,016	41.8 (38.3-45.4)	902			●

^{*}Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN90: Percentage of students who are transgender	1.0 (0.7-1.4)	3,003	4.0 (2.8-5.6)	898		●	
QN91: Percentage of students who never or rarely wore a helmet when they rode a bicycle or used rollerblades or a skateboard (among students who rode a bicycle or used rollerblades or a skateboard during the 12 months before the survey)	88.0 (86.3-89.4)	1,669	84.7 (80.7-87.9)	454			●
QN92: Percentage of students who have been afraid of being beaten up at school (during the 12 months before the survey)	6.8 (5.9-7.8)	3,042	15.0 (12.7-17.7)	911		●	
QN93: Percentage of students who reported that their property had been stolen or deliberately damaged on school property one or more times (such as their car, clothing, or books, during the 12 months before the survey)	17.6 (16.2-19.1)	3,013	27.8 (24.7-31.2)	900		●	
QN94: Percentage of students who have been harassed on school property because someone thought they were gay, lesbian, bisexual, or transgender (during the 12 months before the survey)	6.5 (5.6-7.5)	3,022	21.3 (18.4-24.4)	897		●	
QN95: Percentage of students who have been a member of a gang or crew (during the 12 months before the survey)	17.1 (15.7-18.7)	3,016	17.5 (14.9-20.5)	868			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

District of Columbia (Including Charter Schools) High School Survey Risk Behaviors and Sex of Sexual Contacts Report

Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN96: Percentage of students who have seen or heard people where they live be violent and abusive (including serious hitting, shouting, throwing items, yelling, or name calling, but not 'play fighting,' during the 12 months before the survey)	34.3 (32.5-36.2)	2,999	41.4 (37.8-45.1)	857		●	
QN97: Percentage of students who have ever bullied someone else on school property (during the 12 months before the survey)	8.5 (7.5-9.6)	3,010	13.2 (10.8-15.9)	870		●	
QN98: Percentage of students who think their close friends would strongly approve or approve them having one or more drinks of alcohol nearly every day	12.8 (11.5-14.2)	2,975	19.5 (16.7-22.7)	889		●	
QN99: Percentage of students who think their close friends would strongly approve or approve them using marijuana once a month or more	38.7 (36.7-40.6)	2,994	49.1 (45.5-52.8)	895		●	
QN100: Percentage of students who currently used synthetic marijuana (also called "K2," "Spice," "fake weed," "King Kong," "Yucatan Fire," "Skunk," or "Moon Rocks," one or more times during the 30 days before the survey)	2.4 (1.8-3.1)	2,957	9.0 (7.2-11.3)	853		●	
QN101: Percentage of students who currently used illegal drugs (one or more times during the 30 days before the survey)	16.5 (15.1-18.0)	2,960	27.8 (24.7-31.1)	851		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN102: Percentage of students who reported their partners were three or more years older than themselves the last time they had sexual intercourse (among students who have had sexual intercourse)	8.6 (7.4-9.9)	2,334	12.6 (9.8-15.9)	602		●	
QN103: Percentage of students who have had oral sex with one or more people during their life	62.2 (60.3-64.1)	3,001	72.0 (68.7-75.1)	893		●	
QN104: Percentage of students who reported that they or their partner had not ever used a female condom when they had sexual intercourse during their life (among students who have had sexual intercourse)	86.0 (84.4-87.5)	2,270	82.1 (78.4-85.3)	567	●		
QN105: Percentage of students who have been pregnant or gotten someone pregnant (one or more times)	8.5 (7.4-9.7)	2,941	11.2 (9.0-13.7)	846		●	
QN107: Percentage of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight (also called fasting, during the 30 days before the survey)	12.5 (11.2-13.9)	2,844	20.8 (17.9-24.1)	816		●	
QN108: Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight (during the 30 days before the survey)	6.4 (5.5-7.4)	2,792	14.6 (12.1-17.6)	795		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN109: Percentage of students who ate at a fast food chain or carry out restaurant (at least one day during the 7 days before the survey)	82.4 (80.7-83.9)	2,727	79.8 (76.8-82.6)	833			●
QN110: Percentage of students who most of the time or always went hungry because there was not enough food in their home (during the 30 days before the survey)	4.2 (3.5-5.2)	2,730	9.6 (7.5-12.2)	823		●	
QN111: Percentage of students who reported that none of their classroom teachers provide short physical activity breaks during regular class time (not counting their physical education teacher)	81.5 (79.9-83.1)	2,612	74.9 (71.3-78.2)	756	●		
QN112: Percentage of students who have not ever been tested or are not sure if they have ever been tested for a sexually transmitted disease (STD)	60.4 (58.4-62.4)	2,684	58.5 (54.6-62.2)	793			●
QN113: Percentage of students who reported that there is not at least one teacher or other adult in this school that they can talk to if they have a problem or they were not sure if there is at least one teacher or other adult in this school that they can talk to if they have a problem	33.6 (31.7-35.6)	2,688	35.8 (32.2-39.7)	800			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS
District of Columbia (Including Charter Schools) High School Survey
Risk Behaviors and Sex of Sexual Contacts Report

Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes [†]	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN114: Percentage of students who have not talked with at least one of their parents or guardians about the dangers of tobacco, alcohol, or drug use (during the 12 months before the survey)	52.6 (50.5-54.7)	2,626	49.7 (45.8-53.7)	760			●
QN115: Percentage of students who usually slept in the home of a friend, family member, or other person because they had to leave their home or their parent or guardian cannot afford housing (during the 30 days before the survey)	5.5 (4.6-6.6)	2,697	8.2 (6.3-10.7)	823		●	
QN116: Percentage of students who have ever slept away from their parents or guardians because they were kicked out, ran away, or were abandoned (during the 30 days before the survey)	11.0 (9.7-12.4)	2,629	15.6 (13.0-18.6)	769		●	
QN117: Percentage of students who have ever been given money, a place to stay, food, or something else of value in exchange for sex	6.3 (5.3-7.4)	2,619	11.8 (9.5-14.6)	755		●	
QN118: Percentage of students who sometimes, rarely, or never get the kind of help they need (among students who report having felt sad, empty, hopeless, angry, or anxious)	73.2 (70.9-75.4)	1,680	74.3 (70.2-78.1)	581			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.