

FIRST AID SUPPLIES CHECKLIST

Title 5A DCMR Chapter1, 150.4 and 150.5 – For every 25 children, each facility shall maintain a first aid kit that includes, but is not limited to the following supplies:

A current edition of the first aid text published by the	Telephone number(s) of the local Poison Control
American Academy of Pediatrics, the American Red	Center
Cross, or an equivalent community first aid guide	
□ One roll of one-half inch (½ in.) non-allergenic	□ One roll of two-inch (2 in.) gauze roller
adhesive tape	bandage
□ 10 individually wrapped sterile gauze squares in	□ 25 adhesive compresses, such as
assorted sizes	band-aids, in assorted sizes
Three clean cotton towels or sheeting pieces,	□ One pair of scissors
approximately twenty-four by thirty-six inches (24 in. x	
36 in.) each	
□ Safety pins in assorted sizes	One working flashlight
□ One non-mercury, non-glass thermometer	One measuring tablespoon or dosing spoon
• One pair of tweezers	\Box One-third cup (1/3 c.) of powdered milk for dental
	first aid (for mixing to make a liquid solution)
Rubbing alcohol and alcohol swabs	Cotton balls
□ One ice pack or gel pack	Liquid sanitizer
□ Sanitary soap	Disposable, non-absorbent latex free, or non-
	powdered latex free gloves
□ All items needed for disposal of blood-borne pathogens	Eye patch or dressing
Pen or pencil and notepad	U Wipes
U Whistle	□ One-way valves for infants (if served), young
	children, and adults

- All staff members must possess a current and valid certification in First Aid and CPR.
- Two staff members at the premises are readily available to administer First Aid and CPR (exception: six children or fewer with only two children under the age of two).
- All first aid equipment and supplies must be maintained clean and in a sufficient quantity based on the size of the facility, the ages and developmental abilities of enrolled children, and the facility's program of activities.
- All first aid equipment and supplies must be maintained in a designated location readily available to staff and inaccessible to children.
- Inspection of first aid supplies should occur weekly and be replenished (e.g., removal and replacing sterile supplies if opened, used, damaged, or expired).